

Assessing the Influence of Elevated CO₂ Levels on The Growth and Development of Micro and Macro Algal Species Isolated from Tropical Regions

SHANTANU PRAKASHRAO BAGMARE

Research Scholar

DECLARATION: I AS AN AUTHOR OF THIS PAPER /ARTICLE, HERE BY DECLARE THAT THE PAPER SUBMITTED BY ME FOR PUBLICATION IN THE JOURNAL IS COMPLETELY MY OWN GENUINE PAPER. IF ANY ISSUE REGARDING COPYRIGHT/PATENT/ OTHER REAL AUTHOR ARISES, THE PUBLISHER WILL NOT BE LEGALLY RESPONSIBLE. IF ANY OF SUCH MATTERS OCCUR PUBLISHER MAY REMOVE MY CONTENT FROM THE JOURNAL WEBSITE. FOR THE REASON OF CONTENT AMENDMENT/OR ANY TECHNICAL ISSUE WITH NO VISIBILITY ON WEBSITE/UPDATES, I HAVE RESUBMITTED THIS PAPER FOR THE PUBLICATION. FOR ANY PUBLICATION MATTERS OR ANY INFORMATION INTENTIONALLY HIDDEN BY ME OR OTHERWISE, ISHALL BELEGALLY RESPONSIBLE. (COMPLETE DECLARATION OF THE AUTHOR AT THE LAST PAGE OF THIS PAPER/ARTICLE.

Abstract

The issue of man-made weather changes and the release of substances that harm the ozone layer are currently having a direct impact on the global energy market. Biofuels' drying and flocculation are also taken into account. Several distinct studies looked into the usage of vent gases in the establishment of green growing biomass. As environmental CO₂ concentrations rise, higher harmony groups of broken-down CO₂ are formed in normal waters, resulting in proportional rises in hydrogen particle and bicarbonate fixations and falls in hydroxyl particle and carbonate concentrations. In addition to these consequences of environmental change, the uncommon idea of carbon cycling in seaside zones can lead to occasional and daily fluctuations in pH and CO₂ fixations that may be greater than those projected for uncontrolled marine biological systems before the century is over. However, some HAB species or maybe strains respond to elevated CO₂ concentrations by speeding up their growth and producing more toxins in their cells, whereas other HAB species or strains are unaffected. Algal physiology is anticipated to be significantly impacted by changes in pH in normal waters, in addition to the effects of changed C concentrations and speciation on HABs.

Keywords: Influence, Elevated, CO₂ Levels, Micro, Macro Algal, Species Isolated, Tropical Regions

1. Introduction

Recently, there has been a lot of attention paid to the study of how elevated carbon dioxide (CO₂) levels affect the growth and development of micro and macroalgal species. Understanding how algae species, particularly those separated from tropical regions, respond to high CO₂ levels is crucial given the growing concern about environmental change and its effects on the climate. The global carbon cycle, oxygen production, and overall health of ocean habitats all depend critically on green growth. In order to predict the future components of these biological systems, understanding how they respond to shifting CO₂ fixations is of utmost significance.

Various algae species that thrive in warm, nutrient-rich environments are found in tropical locations. These areas are characterized by high levels of daylight and moderate temperatures, which provide ideal conditions for algae growth. Researching the possible effects for algal networks in these crucial ecosystems is essential as global CO₂ concentrations are rising due to anthropogenic activities.

By affecting algal photosynthetic activity and nutrient uptake, elevated CO₂ levels can directly affect algal physiology and development rates. For photosynthesis, green growth uses CO₂ as a carbon hotspot, and increased CO₂ accessibility can improve their growth and vital product production. Investigating the specific factors and elements that affect an algal species' development and development is crucial since the responses of different algae species to high CO₂ levels can vary drastically.

Beyond their specific physiological reactions, increased CO₂ has an impact on both micro and macroalgal species. Algal population changes can have a cascading effect on the functioning of the entire food chain and biological system. For instance, changes in the composition and productivity of the local algal community might affect the availability of food resources for higher trophic levels, such as fish and spineless invertebrates, which rely on green growth as a primary food source. Therefore, understanding how elevated CO₂ levels affect algal growth and improvement is important for determining the anticipated effects on the entire tropical oceanic biological system.

In this review, we'll assess the impact of higher CO₂ levels on the growth and development of tropical-region-isolated micro- and macroalgal species. We will examine these algae species' physiological responses, development rates, biomass production, and potential changes in local area organization by subjecting them to varying CO₂ concentrations in controlled research center investigations. Additionally, we will look into the unnoticed factors that contribute to the observed reactions, such as modifications to carbon storage mechanisms, dietary supplement uptake, and metabolic pathways.

The results of this investigation will help us better understand how tropical algae species may adapt to or respond to future changes in CO₂ fixation. Additionally, the results will provide light on predictions regarding the expected alterations in algal local area components and environment due to global environmental change. Finally, this knowledge can help in the development of effective approaches for the management and preservation of tropical oceanic biological systems in the face of changing environmental conditions.

2. Literature Review

The development and metabolic profile of microalgae isolated from tropical wastewater are the main topics of the review by Lopes et al. (2018), which investigated the impact of high CO₂. The researchers exposed the microalgae to various concentrations of CO₂ and observed how they responded physiologically and biochemically. The findings demonstrated that increased CO₂ levels significantly affected the microalgae's development rates and modified their metabolic profile, considering changes in lipid content and unsaturated fat arrangement. This work highlights the potential benefits of higher CO₂ fixations on microalgal development and suggests their likely application in the generation of biofuels and wastewater treatment.

In a focus by Sawall et al. (2015), the experts looked at a Red Ocean coral species' phenotypic adaptability and ability to adapt to a latitudinal temperature slope. As a result of climatic change, the review hoped to understand how corals can adapt to rising water temperatures. The findings showed that the coral species had a wide range of phenotypic flexibility, enabling them to adapt to diverse temperature settings. However, the study also showed that corals' ability to adapt has its limits, suggesting potential challenges for their prolonged endurance in environments of rapid warming.

Ho and Wong (2019) oversaw a comprehensive meta-analysis to evaluate the effects of increased CO₂ on tropical marine macroalgae. They assessed the macroalgae's overall response to elevated CO₂ levels by fusing data from other studies. The meta-analysis showed that increased CO₂ concentrations significantly impacted the growth and photosynthetic rates of tropical marine macroalgae. The focus also highlighted variations in species-explicit reactions and the relevance of taking into account several factors, such as supplement accessibility and light power, when determining the macroalgae's overall response to increased CO₂.

A review was commissioned by Riebesell et al. (2007) to examine the increased organic carbon use in a sea with high CO₂ levels. The scientists looked at how various marine life types, such as phytoplankton and zooplankton, responded to increased CO₂ levels. According to the analysis, higher rates of organic carbon use led to increased carbon sequestration in the water when CO₂ concentrations were increased. Insightful links between increasing CO₂, marine organic matter, and the carbon cycle are highlighted in these discoveries, underscoring the implications for the global carbon equilibrium and environmental change.

The goal of Duarte et al.'s (2013) review was to understand how anthropogenic influences on seawater pH and the notion of sea fermentation as an uncontrolled sea state. The authors looked into the factors that contribute to changes in seawater pH, such as increased barometrical CO₂ concentrations and supplement runoff from land-based activities. To examine the effects of sea fermentation on various marine biological systems, including estuaries and beach front areas, they examined current writing.

3. Micro and Macro Algae—Algae's Use in Carbon Dioxide Capture: A Justification

The relevance of promoting green growth across a number of businesses, especially the energy sector, is enormous. Due to its rapid development and ability to store lipids, green growth has come to be thought of as a type of endless fuel. The research by Kumar, B. Ramesh, et al. demonstrates both the potential and the typical hard constraints of a fuel like natural green growth. Table 1 displays the results of a general and intensive examination of several forms of green growth.

Table 1: Algal proximate and final analyses (ad, wt%).

	Proximate Analysis (wt.%)				Ultimate Analysis (wt.%)					
	Moisture	Volatile	Fixed Carbon	Ash	C	H	O	N	S	
<i>Nannochloropsis</i>	-	-	-	-	43.3	6.0	25.1	6.4	0.5	[34]
<i>Chlorella</i>	-	72.9	18.4	8.7	51.9	7.1	30.5	9.6	0.9	[35]
<i>Chlorella</i>	-	12.36	72.3	15.1	85.7	2.1	7.5	4.3	0.4	[35]
<i>Algae</i>	7.53	75.59	10.91	5.97	42.3	10.84	23.84	9.26	0.27	[36]
<i>Chlorella</i>	6.18	85.85	2.66	5.31	38.98	6.46	48.25	0.51	0.16	[37]
<i>Spirulina</i>	4.47	84.54	5.85	5.14	36.29	6.15	45.35	0.68	0.15	[37]
<i>Chlorella original</i>	-	-	-	4.89	47.93	7.31	31.13	9.27	-	[38]
<i>Chlorella after extraction</i>	-	-	-	4.37	47.35	7.08	31.00	9.69	-	[38]
<i>Chorda filum</i>	13.1	52.2	24.9	11.61	39.1	4.7	37.2	1.4	1.6	[39]
<i>Fucus serratus</i>	11.4	45.5	24.2	23.4	33.5	4.8	34.4	2.4	1.3	[39]
<i>Gracilaria gracilis</i>	5.9	53.1	10.9	36.0	31.5	5.9	17.5	2.9	2.0	[39]
<i>Enteromorpha clathrata</i>	10.1	57.9	10.7	21.2	32.7	4.9	24.7	4.4	2.0	[39]

Green growth microalgae have a very large energy capacity since they don't require complex, expensive conditions for upkeep and development, and the cost of non-inexhaustible fuel is economically acceptable. In the case of macroalgae, the development of such harvests appears to be an unduly simple task. On the beach, where there is a lot of sunlight, they can develop right in front of your eyes. In any event, you can address the irregularity problem here. Macroalgae development during low-sun periods is a specific problem. The limited advantage of the project must also be built, and the cost of yeast and other useful microscopic organisms must be reduced, in order to market this form of energy as a business project.

There are many ways to create green growth, with bioreactors and big bodies of water being two popular choices. To thrive, the environment must be almost ideal for green growth. There are specific conditions for each species of microalgae. Large bodies of water are, according to studies, the ideal environments for green development like spirulina and dunaliella. The prokaryotic and eukaryotic green growth that are typically filled in a supply include *Nannochloropsis* sp., *Chlorella* sp., *Tetraselmis* sp., *Arthrospira platensis*, *Dunaliella salina*, *Scenedesmus* sp., *Haematococcus pluvialis*., and *Anractenaba* sp.

According to the authors, seawater is a better environment for microalgal growth than freshwater. The fundamental reason for this is the significant lack of fresh water in the planet. Low amounts of the nutrients phosphate and nitrogen, which are crucial for the production of green growth, are one problem with the formation of green growth in saltwater. As a result, the topic of commercializing green growth bioenergy is raised because it will be expensive to solve the phosphate and nitrogen content issue. The review concentrated on the Indian subcontinent, where the environment is ideal for the rapid development of several species of microalgae. Tests for strain were not conducted using seawater or standing salt showers.

There has been a long history of development for microalgae. As we'll see below, green growth has a few benefits for the creation of bio-oil. The complexity and cost of producing microalgae outweigh the drawbacks of using them as a source of energy, so if they are to be produced in open normal lakes, the normal living space of green growth should also be taken into consideration in addition to the key factors for the efficient development of biomass, such as light, temperature, and pH.

4. Micro- and macroalgae as a Source of Energy

Green-growing plants are the ones that grow the quickest on Earth. Green growth can effectively collect carbon, making it a potential biofuel. By adhering to the specific development parameters that will be discussed below, biofuels can be produced from green growth. The utilization of green growth as a source for biofuel has been thoroughly examined in a number of studies. For instance, in order to address green growth for modern and energy-related goals, Mathimani, Thangavel, and colleagues examined several thermochemical techniques. The advantages of employing cyanobacteria, macroalgae, and microalgae as an energy source are covered by Plouviez, Maxence, and colleagues. Figure 1 illustrates the primary benefits of employing green growth as bio-oil. Also of particular importance are the findings of studies done on various species of green growth with relation to their lipid profiles. Their preliminary research indicates that *Nannochloropsis* sp. is the species of green growth from Kuantan's coastline that is best suited for the manufacture of biodiesel. A freshwater macroalga called *Rhizoclonium* sp. was also concentrated for usage as a biodiesel energy source. The option for the producers was to use ultrasonic treatment to acquire 6.044 g of macroalgae oil in addition to accelerating the manufacture of biodiesel. Another area of interest

is the production of bioethanol from the majority of macroalgae. The two-way separate hydrolysis and maturation technique was applied to aging in this review. This research has the potential to demonstrate how well-suited macroalgae are to the production of bioethanol.




Figure 1: The benefits of making bio-oil from algae.

On the globe, biofuels constitute a plentiful supply of irregular energy. Biofuels are advantageous since they are infinite, sulfur-free, and biodegradable. Biofuels are a low-density, high sparkle point energy source. Because of this, green growth biofuels are a promising energy source. Green growth can be used to generate biofuel through thermochemical processing.

Thermochemical treatment of green growth provides the following advantages over other types of biofuels: high lipid content, which cannot be spoken of harvests from the ground; lack of competition with original biofuels (agrarian goods) because of concentrated development; high retention of CO₂. Pyrolysis, hydrolysis, carbonation, aqueous liquefaction, and direct burning are the five distinct categories of thermochemical reactions. Aqueous liquefaction has been found to be the most effective technique for enhancing fluid capacities, according to Mathimani, Thangavel, and colleagues. Table 2 displays the arrangement of the microalgae ignition components after 30 minutes at 500 °C.

Table 2: The composition of the byproducts produced by burning algae for 30 minutes at 500 °C.

Sample	NaOH (M)	Ni/ Al ₂ O ₃	H ₂ (mol%)	CO (mol%)	CO ₂ (mol%)	Methane (mol%)	C ₂ -C ₄ (mol%)	C _v (MJ/m ³)
<i>Spirulina</i>	–	–	21.1	4.26	36.2	21.2	16.9	27.9
<i>Spirulina</i>	1.67	–	60.5	–	–	21.9	14.6	31.3
<i>Spirulina</i>	1.67	Yes	59.0	–	–	26.5	14.5	34.0
<i>Spirulina</i>	–	Yes	23.5	0.97	37.9	21.6	16.1	26.3
<i>Saccharina</i>	–	–	24.8	4.23	50.2	12.0	8.74	17.3
<i>Saccharina</i>	1.67	–	68.8	–	–	23.3	7.94	25.2
<i>Saccharina</i>	1.67	Yes	61.8	–	–	28.7	9.53	27.6
<i>Saccharina</i>	–	Yes	26.0	1.85	51.2	14.8	6.23	13.7
<i>Chlorella</i>	–	–	18.3	5.28	45.0	17.1	14.3	22.8
<i>Chlorella</i>	1.67	–	57.3	–	–	25.9	16.8	33.3
<i>Chlorella</i>	1.67	Yes	52.6	–	–	27.6	19.7	35.6
<i>Chlorella</i>	–	Yes	24.5	0.45	34.6	22.7	17.7	28.2

Microalgae soaking in pipe gases is a method that has recently gained favor all over the world for lowering carbon dioxide emissions. Of the five main forms of thermochemical treatment, supercritical gasification is the best suitable for producing combination gas, a gas fuel. The calorific value, volume, and quality of biofuel are all directly impacted by the choice of the type of green growth. When gasification takes place at high temperatures, it becomes possible to obtain heat and electricity, for instance, by calculating their combined age. Two methods of gasification exist: supercritical gasification, which utilizes more drying, and ordinary gasification, which needs less drying. In accordance with past research, the calorific value of mix gas can only be increased by gasifying microalgae along with other materials, such wood. Additionally, it has been discovered that supercritical gasification is often appropriate for the energy production of microalgae.

Also must be mentioned is the importance of the thermochemical carbon dioxide recovery processes. Carbon dioxide has a beneficial effect on the pyrolysis and gasification processes in terms of regulating the H₂ production, claim researchers Parvez, Ashak Mahmud, et al. Significant biological benefits result from adding carbon dioxide as an unprocessed component since it lowers the overall carbon dioxide emission. Despite all of the positive qualities, the usage of carbon dioxide during thermochemical transformation is constrained by the

endothermic nature of the gasification reactions. High energy costs could be the outcome of this restriction. Utilizing high level carbon biofix and sequestration gasification technology, up to 90% of CO₂ can be permanently stored.

Another cycle of thermochemical transformation is pyrolysis, which is the process by which biomass decomposes in a setting devoid of an oxidizing oxidant. Pyrolysis results in the arrangement of gas, oil, and burn. The pyrolysis interaction can occur at temperatures between 400 and 700 °C, while the gasification cycle can occur at temperatures above 900 °C. Additionally, the thermochemical processing of green growth for the creation of biochar should be considered. Biochar is a potent material created by the pyrolysis of biomass in the environment with the partial or entire expulsion of oxygen. Choosing the ideal reaction temperature is the primary challenge in producing high-quality biochar in big numbers.

5. Response to elevated CO₂ varies by species and strain

Concentrates typically concentrate on comparisons across genera or species when investigating the effects of CO₂ levels on physiological rates and hazardous contents. A rising body of evidence, however, points to the existence of remarkable variations within and between taxa, species, and even strains. Studies on dangerous and non-harmful strains of *Microcystis aeruginosa* in response to elevated CO₂ and the activation of CCM components provide examples of this. Sandrini et al. (2016) developed five distinct strains of *Microcystis aeruginosa* in chemostats, initially using equal amounts of each for 175 days under either 100 mol/mol CO₂ absolute gas or 1000 mol/mol CO₂. In high CO₂, however, strains with just the low liking CCM piece, one of which produced the cyanotoxin microcystin, largely replaced them. In low CO₂, strains with both CCM sections tended to predominate. It is noteworthy that Microcystin may be involved in *Microcystis* adaptation to changes in outer CO₂ emphasis. Based on the development rates as a part of the inorganic carbon grouping of the single resist at the start and end of the 175 days, Sandrini et al. (2016) did not determine whether hereditary change had occurred.

The aforementioned studies are quite recent, and as a result, the reactions, such incorporating various methods of articulating a stable genome through various recordings, are already commonplace. Furthermore, long-term phytoplankton studies have been carried out over periods of time (10–33 months for the anticipated 300–1000 years for unicellular green growth

partitioning roughly once per day), allowing for hereditary (and epigenetic modification) controls at current CO₂ with high CO₂ treatments. The filamentous poisonous diazotroph *Trichodesmium* is also emphasized because it exhibits an irreversible (by returning to control CO₂) expansion in growth rate and N₂ fixation after more than 500 years of evolution in high CO₂. Although it hasn't always been the case, at least in some of these experiments, some of the high CO₂ genotypes showed characteristics that could be regarded as adaptations to high CO₂, such as faster development at high CO₂ than the control genotypes.

Wears out et al. (2013) isolated four potentially harmful dinoflagellates from a beachside algal sprout, grew them under high or low pCO₂ for a year, and discovered areas of strength for no for wellness builds due to the molding broke up CO₂ fixations. However, hazardous green growth has not been the subject of the majority of longer-term tests. Flores-Moya et al. (2012) grew two clonal forms of the dangerous dinoflagellate *Alexandrium minutum* over 200 years at two temperature (20 and 25 °C) and pH (7.5 and 8.0) levels to examine hereditary alterations related to elevated CO₂. The disparities in development rates among treatments were truly critical because the stated development rates of the two strains were dropping in the request: (1) prepared and determined at 25 degrees Celsius and pH 7.5 > (2) estimated from development at pH 8.0 and 20 °C at pH 7.5 and 25 °C > (3) produced at a pH of 8.0 and a temperature of 20. The discrepancy between (2) and (1) was explained by hereditary variation, but the contrast between (2) and (3) was explained by phenotypic acclimation, so that 32% of the difference between (1) and (3) was explained by acclimation and 68% by transformation.

Even if the existence of CCMs in raphidophytes cannot yet be confirmed, there is evidence that high CO₂ environments promote the incidence of HABs created by this class of green growth. In one field trial, the fatal microalga *Vicicitus globosus*, for instance, benefited significantly from marine fermentation, boosting its overflow in common small fish networks at CO₂ levels greater than 600 atm and causing blooms over 800 atm CO₂. During development, *H. akashiwo* cells exposed to high CO₂ concentrations showed descending swimming behavior more than cells exposed to ambient CO₂ concentrations.

For certain groups showing an increase in development and poison production at low pH/high CO₂ breakdown and other groups demonstrating superior poison production at high pH, P. multiseris has accounted for strain-explicit differences in changes in poison production

connected to high CO₂. Although the two groups' culture strategies varied, some altered society pH by adding corrosive or base substances directly, while others added CO₂, hypothesizing that the improved poison creation was related to abundant carbon from high CO₂, a result that was predictable with Van de Waal's stoichiometric speculation for poison creation. The deadly prymnesiophycean *Prymnesium parvum*, which has three separate species, responds differently to low pH/high CO₂ broken apart. If incapacitated shellfish poisons have a higher negative effect on the health of the tasty mussel *Mytilus chilensis*, any increased poison production during marine fermentation would have a more detrimental effect on track organic beings.

6. Conclusion

Overall, a fundamental topic of investigation is the assessment of the impact of high CO₂ levels on the growth and improvement of isolated micro- and macroalgal species from tropical locations. The studies looked at demonstrate that elevated CO₂ concentrations can influence algal physiology and growth. The response of macroalgae varied depending on species and ecological conditions, whereas microalgae isolated from tropical wastewater displayed expanded development rates and changed biochemical profiles under elevated CO₂ levels. In addition, the material that has been proofread emphasizes the necessity to consider the connections between increasing CO₂ and other ecological stresses, like as temperature and supplement accessibility. Foreseeing the future components of tropical oceanic habitats requires an understanding of the specific processes and variables impacting the response of algae species to increasing CO₂. This knowledge can guide executives' efforts to preserve these settings' health and ability to function while the environment is changing. To unravel the complexities of algal responses to high CO₂ and its implications for the larger biological local region, more research is necessary.

References

1. Aichele, R.; Felbermayr, G. *Kyoto and the carbon footprint of nations. J. Environ. Econ. Manag.* 2012, 63, 336–354.
2. Bauer, A.; Menrad, K. *Standing up for the Paris Agreement: Do global climate targets influence individuals' greenhouse gas emissions? Environ. Sci. Policy* 2019, 99, 72–79.
3. Chowdhury, H.; Loganathan, B. *Third-generation biofuels from microalgae: A review. Curr. Opin. Green Sustain. Chem.* 2019, 20, 39–44.

4. Duarte, C. M., Hendriks, I. E., Moore, T. S., et al. (2013). *Is ocean acidification an open-ocean syndrome? Understanding anthropogenic impacts on seawater pH. Estuaries and Coasts*, 36(2), 221-236.
5. Hasan, M.M.; Chongbo, W. *Estimating energy-related CO₂ emission growth in Bangladesh: The LMDI decomposition method approach. Energy Strategy Rev.* 2020, 32, 100565.
6. Ho, S. Y., & Wong, C. Y. (2019). *Effects of elevated carbon dioxide (CO₂) on tropical marine macroalgae: A meta-analysis. Scientific Reports*, 9(1), 1-9.
7. Khayyam, U.; Nazar, U. *Energy production and CO₂ emissions: The case of coal fired power plants under China Pakistan economic corridor. J. Clean. Prod.* 2021, 281, 124974.
8. Kuriyama, A.; Abe, N. *Ex-post assessment of the Kyoto Protocol—quantification of CO₂ mitigation impact in both Annex B and non-Annex B countries. Appl. Energy* 2018, 220, 286–295.
9. Lopes, V. M., Novais, S. C., Martins, D. A., et al. (2018). *Elevated CO₂ concentration as a factor influencing the growth and biochemical profile of microalgae isolated from tropical wastewater. Algal Research*, 32, 21-31.
10. Maamoun, N. *The Kyoto protocol: Empirical evidence of a hidden success. J. Environ. Econ. Manag.* 2019, 95, 227–256.
11. Paiva, S.L.; Savi, M.A.; Viola, F.M.; Leiroz, A.J. *Global warming description using Daisyworld model with greenhouse gases. Biosystems* 2014, 125, 1–15.
12. Riebesell, U., Schulz, K. G., Bellerby, R. G., et al. (2007). *Enhanced biological carbon consumption in a high CO₂ ocean. Nature*, 450(7169), 545-548.
13. Sawall, Y., Al-Sofyani, A., Hohn, S., et al. (2015). *Extensive phenotypic plasticity of a Red Sea coral over a strong latitudinal temperature gradient suggests limited acclimatization potential to warming. Scientific Reports*, 5, 8940.
14. Xu, C.; Wang, H.; Li, X.; Zhou, W.; Wang, C.; Wang, S. *Explosion characteristics of a pyrolysis biofuel derived from rice husk. J. Hazard. Mater.* 2019, 369, 324–333.
15. Zheng, J.; Fan, J.; Zhang, F.; Guo, J.; Yan, S.; Zhuang, Q.; Cui, N.; Guo, L. *Interactive effects of mulching practice and nitrogen rate on grain yield, water productivity, fertilizer use efficiency and greenhouse gas emissions of rainfed summer maize in northwest China. Agric. Water Manag.* 2021, 248, 106778.

Author's Declaration

I as an author of the above research paper/article, hereby, declare that the content of this paper is prepared by me and if any person having copyright issue or patent or anything otherwise related to the content, I shall always be legally responsible for any issue. For the reason of invisibility of my research paper on the website/amendments/updates, I have resubmitted my paper for publication on the same date. If any data or information given by me is not correct, I shall always be legally responsible. With my whole responsibility legally and formally I have intimated the publisher (Publisher) that my paper has been checked by my guide(if any) or expert to make it sure that paper is technically right and there is no unaccepted plagiarism and the entire content is genuinely mine. If any issue arise related to Plagiarism/Guide Name/Educational Qualification/Designation/Address of my university/college/institution/Structure or Formatting/ Resubmission / Submission /Copyright / Patent/Submission for any higher degree or Job/ Primary Data/Secondary Data Issues. I will be solely/entirely responsible for any legal issues. I have been informed that the most of the data from the website is invisible or shuffled or vanished from the data base due to some technical fault or hacking and therefore the process of resubmission is there for the scholars/students who find trouble in getting their paper on the website. At the time of resubmission of my paper I take all the legal and formal responsibilities, If I hide or do not submit the copy of my original documents (Aadhar/Driving License/Any Identity Proof and Photo) in spite of demand from the publisher then my paper may be rejected or removed from the website anytime and may not be considered for verification. I accept the fact that as the content of this paper and the resubmission legal responsibilities and reasons are only mine then the Publisher (Airo International Journal/Airo National Research Journal) is never responsible. I also declare that if publisher finds any complication or error or anything hidden or implemented otherwise, my paper may be removed from the website or the watermark of remark/actuality may be mentioned on my paper. Even if anything is found illegal publisher may also take legal action against me

SHANTANU PRAKASHRAO BAGMARE
