

BIOREMEDIATION OF SELECTED SOLID WASTES BY CONTROL
CULTIVATION OF OYSTER MUSHROOM, *Pleurotus florida*

¹Oinam Jennifer,² Laishram Joymati and ³A.Subhashini

¹ Research Scholar, Dept. of Biotechnology, PAHER University, Udaipur, Rajasthan

² Asst. Professor, P.G. Dept. of Zoology, D.M.College of Science, Imphal, Manipur.

³.Research Scholar, P.G. Dept. of Zoology, D.M.College of Science, Imphal, Manipur

Abstract

Mushrooms are a source of nutrition particularly in respect of proteins and vitamins. It has been rightly called “vegetable meat”. Apart from being potential protein and energy source, mushrooms are famous for their medicinal values. In addition to their fascination for the gourmet, they are nature’s most active agents in the disposal of forest’s waste materials. Not only in disposing timber, but also helps in converting dead plants into available food. They are efficient agents, fitting into the nutrient cycle of farms and forests. As they utilize agricultural by products and wastes as their substrate, mushroom cultivation is an eco friendly activity. In a country like India where vegetarians dominate, every attempt should be made to popularize a vegetable protein source like mushroom.

A study on the cultivation of oyster mushroom, *Pleurotus florida* was undertaken. It was found to grow on various solid wastes viz. paddy straw, rice husk, saw dust, water hyacinth and a mixture of all the selected substrates. The oyster mushroom cultivated on the paddy straw gives the maximum yield with high content of protein, carbohydrate and lipid.

Key words: eco- friendly, *Pleurotus florida*, nutritional contents

INTRODUCTION:

According to Chang (1991), “ the mushroom is a macrofungus with distinctive fruiting body which can be either epigeous or hypogeous and large enough to be seen with naked eye and to be picked by hand.” Mushroom farms may be use as disposal sites for agricultural waste products. Mushrooms are found to possess the two basic abilities of biodegradation and bioaccumulation for the bioremediation of wastes. Mushroom cultivation serves as the most efficient and economically viable biotechnology for the conversion of long cellulose waste materials into high quality protein food (Dike *et.al.*,2011). The protein-extracting ability of oyster mushrooms makes them a very

popular alternative energy source for dieting individuals (Shah,2004). Mushrooms are able to convert substrate mass into edible mushroom biomass often exceeding 100% biological efficiency.

A study on the cultivation of *Pleurotus florida* was undertaken. It was cultivated on five selected solid wastes *viz* .Paddy straw, Rice husks,water hyacinth, Saw dust and a mixture of all the selected four substrates. The cultivation was carried out in polythene bags (containing 500gm paddy straw on dry weight basis per bag) using grain spawn as well as the nutritional contents like the protein, lipid and carbohydrate were estimated.

MATERIALS AND METHOD:

Cultivation of the species, *Pleurotus florida*

Materials :

1. Mushroom Spawn
2. Paddy straw, rice husks, water hyacinth and saw dust for making bed substrates :500gm/bag.
3. Perforated transparent polythene bags: 30x 40cm

4. Spawn:100gms/bag
5. Sprayer:1 litre capacity
6. Containers for sterilization.
7. Binding threads.

Methodology:

- Spawn preparation:

The primary inoculum was prepared from the fresh fruiting body of the mushroom through **tissue culture** method and was maintained and multiplied by sub culturing on sterilized PDA medium in petri dishes and test tubes. The spawn was prepared on coarse paddy grains. The grains were half boiled and filled in transparent polythene bags, which were then sterilized in an autoclave at 15psi for 30 minutes. The inoculation was made on the following day under aseptic conditions and incubated at room temperature till the grains were covered with white mycelial growth.

- Substrate preparation:

Pleurotus florida was grown on the selected substrates viz. paddy straw, rice husks, water hyacinth, saw dust and mixture of the four selected

substrates. The substrate paddy straw was obtained from a nearby farmland, rice husk from a rice mill, saw dust from a saw mill and water hyacinth from a nearby pond.

- Soaking:

The substrates were chopped into 6-8cm pieces and soaked in fresh water separately for a minimum of 20 hours and a maximum of an overnight.

- Sterilization and spawning:

The substrates were boiled for at least 30 minutes and were let to cool down at room temperature and the excess water was strained. The substrates were then filled in the perforated polythene bags layer by layer sandwiching the spawn grain respectively. The open ends were then bound by the binding threads and kept at a temperature maintained at

about 30°C and well protected from pests.

- **Cropping and harvest:**

After the spawn run has completed and the bags were fully impregnated with white mycelia, the polythene bags were removed carefully. The

open blocks were kept on racks about 20cm apart. Water was sprayed regularly to keep it moist. Within 3-4 days of the spawn run pin-heads were seen and within another 3-4 days mushrooms were ready for harvest.

ANALYSIS FOR NUTRITIONAL CONTENTS OF THE MUSHROOM HARVESTED:

The mushroom harvested from the various substrates were analysed for the following-

I) Protein content (Lowry et.al., 1951)

The protein content of the mushroom was estimated by the method described by Lowry et.al.,(1951)

II) Carbohydrate content (Hedge and Hofreiter,1962)

The carbohydrate content of the mushroom was estimated by the method described by

Hedge and Hofreiter,(1962)

III) Total Lipid content (Folch,1954)

The total lipid content of the mushroom was estimated by using the method described by Folch,1954.

IV) Biological Efficiency:

The biological efficiency was calculated as a percentage of the yield of the fresh mushroom to the dry weight of the substrates.

RESULTS:

i) Sporophore initiation period and first (in days) of *Pleurotus florida*

Sl.no.	Substrates	Sporophore initiation (in days)	First flush (in days)
1.	Paddy straw	16	21
2.	Water hyacinth	21	28
3.	Rice Husk	18	21
4.	Saw dust	23	30
5.	Mixed substrate	20	25

Different substrates showed different period (in days) for sporophore initiation. A minimum period of 16 days was taken by paddy straw for sporophore initiation followed by rice husk (18 days), mixed substrate (20 days), water hyacinth (21days) and a maximum period of 23 days was recorded in saw dust substrate.

Regarding the first flush, paddy straw and rice husk gave first flush in a minimum of 21days followed by mixed substrate in 25 days, water hyacinth in 28 days and a maximum of 30 days by the saw dust substrate.

ii) Total yield (in grams) of *P.florida* on different substrates in total number of days.

Sl.no.	Substrates	Total number of days taken	Total yield* (gm/500gm)

1.	Paddy straw	42	398.78
2.	Water hyacinth	54	256.41
3.	Rice Husk	60	281.81
4.	Saw dust	66	170.62
5.	Mixed substrate	49	284.84

* Mean of replicates

The order of efficiency of total production in 500 gms substrate is-
 paddy straw (398.78gms/500gms) > mixed substrate (284.83gms) > rice husk (281.81gms) > water hyacinth (256.41gms) > saw dust (170.62gms).

iii) Yield and biological efficiency of *P.florida* grown on different substrates.

Sl.no.	Substrates	Total yield* (gm/500gm)	Biological efficiency(%)
1.	Paddy straw	398.78	79.75
2.	Water hyacinth	256.41	64.10
3.	Rice Husk	281.81	56.36
4.	Saw dust	170.62	34.12
5.	Mixed substrate	284.84	71.21

The order of biological efficiency is paddy straw (79.75%) > mixed substrate (71.21%) > water hyacinth (64.10%) > rice husk (56.36%) > saw dust (34.12%).

iv) Protein ,Carbohydrate and lipid profile of *P.florida* cultivated on different substrates.

Sl.no.	Substrates	Protein (%)	Carbohydrate (%)	Lipid (%)	Caloric value (cal)
1.	Paddy straw	26.2	10.68	0.3	194.67
2.	Water hyacinth	22.5	7.69	0.3	161.47
3.	Rice husk	25.0	8.82	0.1	178.35
4.	Saw dust	20.0	9.97	0.2	155.76
5.	Mixed substrate	25.0	8.95	0.1	178.97

The order of protein (%) of *P.florida* grown on different substrates is paddy straw (26.2%)> rice husk (25.0%)=Mixed substrate(25.0%)> water hyacinth(22.5%) > saw dust(20.0%).

The order of carbohydrate content(%) of *P.florida* grown on different substrates is paddy straw(26.2%)> saw dust(9.97%) >mixed substrate(8.95%)>rice husk(8.82%) >water hyacinth(7.69%)

The order based on lipid content of *P.florida* grown on different substrates is paddy straw(0.3%) = water hyacinth(0.3%)>saw dust(0.2%)>rice husk(0.1%)=mixed substrate(0.1%).

The order based on the caloric value (cal) is paddy straw (194.67) >mixed substrate(178.97) >rice husk(178.35)>water hyacinth(161.47)>saw dust(155.76).

DISCUSSION:

India being an agricultural country, agricultural wastes such as

paddy straw and rice husk are available abundantly which if not properly used will pollute

the environment. Water hyacinth is also considered to be a menace to the aquatic environment as well as in navigation. These wastes can be successfully utilized for mushroom cultivation and it is a way of solid waste management too.

Suitability of paddy straw as bed substrates for oyster mushroom cultivation was reported by Bano and Srivastav (1962) and Block *et al.* (1958). Zadrazil and Schneiderei (1972) too observed that rice husk can be successfully used for the oyster mushroom cultivation. Pani and Mohanty (1998) also reported water hyacinth to be a suitable substrate for oyster mushroom cultivation. Sharma and Jandaik (1981) reported the use of mixed substrates as bedding material with the view that mixed substrates provide a more balanced supply of nutrients to the mushroom. These findings were found to be similar with the present work. Besides, other agro wastes such as saw dust, rice husk, maize cobs, coir pith etc. which are mostly burnt leading to waste of energy

pollution of the environment, can be converted into food, feed and fuel by oyster mushroom cultivation, representing the only economically profitable biotechnological process for the bioconversion of waste plant residues.

In the present study quick sporophore initiation was observed in the case of paddy straw followed by rice husk, mixed substrate, water hyacinth and least in saw dust. The reason being the dependence of the sporophore initiation upon the nature of substrates used as bedding material. Similar reason was stated by Zadrazil (1978). It was found that paddy straw gave the highest first flush followed by water hyacinth. Similar observations were found in the study of Bahram (1989), Jandaik (1989) and Marimuthu *et al.* (1991). Superiority of the paddy straw as a substrate was reported by several workers (Pal and Thapa, 1979; Bahram, 1989; Jandaik, 1989 and Marimuthu *et al.*, 1991). They also reported a maximum biological

efficiency of *P.florida* grown on paddy straw, which was found similar to the results of the present study. The present study showed high level of protein and minimum level of

lipid in the *P.florida* grown on different substrates. The findings of Randive, 2012 support the findings of the present study.

CONCLUSION:

The highest yield, highest biological efficiency, maximum protein, carbohydrate and caloric value were recorded in the *P.florida* grown on paddy straw substrates while the least were seen in the case of saw dust grown *P.florida*. To conclude, paddy straw among

the five substrates used is more suitable for the cultivation of *P.florida*. Though mushroom cultivation, combating malnutrition and protection of the environment is also possible for the successful sustenance of human life and ecosystem.

REFERENCES

1. Bahram, H., 1989. Mushroom Journal,9:97-100.
2. Bano,Z and Srivastav,1962.Studieson the cultivation of *Pleurotus* species,Food Science,11:363-365.
3. Block, S.S., Tsao,G. and Han, L., 1958. Production of mushrooms from saw dust. Journal of Agricultural and Food Chemistry,6: 923-927.
4. Dike K.S.,Amuneke E.H.,and Ogbulie J.N., 2011.J Microbiol.Biotech.Res.,2011, 1(3):1-4.
5. Folch,J.,Lees,M.and Sloane Stanley,G.H.,1957.A simple method for the isolation and purification of total lipids from animal tissues,J.Biol.Chem.,266:497-509.

6. Hedge, J.E. and Hofreiter, B.T., 1962. In: Carbohydrate Chemistry, 17, Academic Press, New York.
7. Lowry, O.H., Rosebrough, N.J., Farr, A.L. and Randa, R.J., 1951. J. Biol. Chem., 193:265.
8. Marimuthu, T., Krishnamoorthy, A.S. and Jayarajan, R., 1991. Glimpses of Mushroom Research in Tamil Nadu Agricultural University, Coimbatore, P.63.
9. Pal, J. and Thapa, C.D., 1979. Cultivation of Dhingri made easy, Indian Journal of Mushrooms, 5:7-20.
10. Pani, B.K. and Mohanty, A.K., 1998. Utilization of water hyacinth as an alternative substrate for oyster mushroom cultivation, Crop Res., 15:294.
11. Shah Z.A., Ashraf M., & Istiaq M. 2004. Comparative Study on Cultivation and Yield performance of Oyster Mushroom (*Pleurotus ostreatus*) on Different Substrates (Wheat Straw, Leaves, Saw dust). Pakistan Journal of Nutrition 3(3):158-160
12. Sharma, A.D., and Jandaik, C.L., 1981, Quality of paddy straw and its effect on spawn run and yield of *Pleurotus sajor-caju*, Indian Journal of Mushroom, pp.84-87.
13. Sonali D. Randive, 2012. Cultivation and study of growth of oyster mushroom on different agricultural wastes substrate and its nutrient analysis: Advances in Applied Science Research, 3(4):1938-1949.
14. Zadrazil, F. and Schneiderei, M., 1972. Die Grundlagen for die Inkultumahme einer bisher

nicht kultivierten *Pleurotus*
Art, *Der Champion*, 12:25-32
15. Zadrazil, F., 1978. Cultivation
of *Pleurotus*. In the *Biology*
and Cultivation of edible

mushrooms, (ed. S.T. and
W.A. Hayes), Academic
Press, New York, pp. 251-
256.

AIRO INTERNATIONAL JOURNAL